



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Triton Alpestris was the form mainly used in this study, and one principle object which the author had in mind was to determine whether the development of the spinal cord took place in a manner similar to that described by His for man. The conclusions support those of His. The mitoses which give rise to the spongioblasts of these authors take place earlier than those which form the neuroblasts—the first form of nerve cells. The supporting substance of the cord is essentially epidermal, therefore, though, in the adult, cells of a different nature may be found imbedded in it. The ganglion cells are of several sizes, and it is the largest ones that develop earliest. Structures which have been described as “granules” and “free nuclei” are, in some cases at least, small ganglion cells. Triton also shows large nerve cells which are the homologues of the “posterior cells” (Freud) in *Petromyzon*. The plates that accompany the paper show several cross sections of the cord, and it is remarkable how closely the early stages resemble the developing cord in man.

Ueber den oberen Kern des Nervus oculomotorius. Dr. L. DARKSCHEWITSCH. Arch. f. Anat. u. Entwicklungsgesch. January, 1889. H. I. and II. Taf. I.

By the study of cross-sections from the region of the anterior corpora quadrigemina in the human fœtus, between the seventh and eighth months, Darkschewitsch makes out a group of cells to which he gives the name superior nucleus of the oculomotorius. The following is taken from his description: There are in this region two columns of cells on each side, their long axis parallel to the aquæduct. The more ventral and caudal group lies nearer the middle line, the more dorsal and cephalic one being laterad of it. The latter group has much smaller cells than the former. In their relations to the oculomotor nerve fibers and the posterior longitudinal bundle, both groups are alike. It is this dorsal and cephalic group, composed of the small cells, which is the “superior nucleus” of our author. For its relations, see the original paper. (Gudden has already described the several cell clusters which form the oculo-motor nucleus in the rabbit, and it may be that a study of this superior nucleus in the adult human brain will make it possible to homologize the subdivisions in man and the rabbit. In the meantime it must be remembered that this “superior nucleus” is classed with the oculo-motor centre solely on the ground of juxtaposition and its relation to the posterior longitudinal bundle and the oculo-motor nerve.—REV.)

Multiple Hirnnervenlähmung nach Basisfractur. Ein Beitrag zur Frage des Verlaufs der Geschmacksnerven. L. BRUNS. Archiv f. Psychiatrie, u. Nervenkrankheiten, Bd. XX, H. 2, 1889.

The patient was a man who had been thrown from a wagon violently on his head; several cranial nerves (from II–VII inclusive) were injured by what was diagnosed as fracture of the basis of the skull. The careful examination showed that in general there existed on the right side, on which there was total paralysis of the facialis, a complete hemiageusia, both at the tip and back of tongue and soft palate, while on the left side, on which the trigeminus was completely paralyzed sensibility to taste was everywhere retained. There was no evidence that the glosso-pharyngeus was injured other than was furnished by the loss of taste. It was further surmised that the trigeminal lesion was intracranial, while that of the facialis was in the Fallopian canal. If the hemiageusia had been confined to the anterior two-thirds of the tongue, the case would have fitted nicely with the theory of Carl, which makes the course of the taste fibers from the glosso-pharyngeus—where they arise—through the ganglion petrosum and by the tympanic nerve to the tympanic plexus, from here the main portion passes by the nervus